

Some Types of HyperNeutrosophic Set (3): Dynamic, Quadripartitioned, Pentapartitioned, Heptapartitioned, m-polar, pp. 178-192, in Takaaki Fujita, Florentin Smarandache: *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond. Fourth volume: HyperUncertain Set (Collected Papers)*. Gallup, NM, United States of America – Guayaquil (Ecuador): NSIA Publishing House, 2025, 314 p.

Chapter 5

Some Types of HyperNeutrosophic Set (3): Dynamic, Quadripartitioned, Pentapartitioned, Heptapartitioned, m-polar

Takaaki Fujita^{1*}, Florentin Smarandache²

^{1*} Independent Researcher, Shinjuku, Shinjuku-ku, Tokyo, Japan. t171d603@gunma-u.ac.jp

² University of New Mexico, Gallup Campus, NM 87301, USA. smarand@unm.edu

Abstract

This paper builds upon the foundation established in [50, 51]. The Neutrosophic Set provides a robust mathematical framework for handling uncertainty, defined by three membership functions: truth, indeterminacy, and falsity. Recent developments have introduced extensions such as the Hyperneutrosophic Set and SuperHyperneutrosophic Set to tackle increasingly complex and multidimensional problems.

In this study, we explore further extensions, including the Dynamic Neutrosophic Set, Quadripartitioned Neutrosophic Set, Pentapartitioned Neutrosophic Set, Heptapartitioned Neutrosophic Set, and m-Polar Neutrosophic Set, to address advanced challenges and applications.

Keywords: Set Theory, SuperhyperNeutrosophic set, Neutrosophic Set, HyperNeutrosophic set

1 Preliminaries and Definitions

This section introduces the key concepts and definitions fundamental to the discussions in this paper. The study employs standard set-theoretic operations and extends them to advanced frameworks. Readers seeking an in-depth understanding of classical set theory are referred to resources such as [25, 68, 71, 72]. For foundational principles and applications of Neutrosophic Sets, the referenced literature provides comprehensive insights.

1.1 Neutrosophic, HyperNeutrosophic, and n-SuperHyperNeutrosophic Sets

Various set-theoretic frameworks have been devised to address uncertainty, vagueness, and imprecision in decision-making. These frameworks include Fuzzy Sets [120–124], Intuitionistic Fuzzy Sets [6–11], Vague Sets [2, 12, 64, 73, 97], Plithogenic Sets [27, 34, 37–39, 47, 48, 60, 109, 111, 112], Soft Sets [65, 74, 79], Hypersoft Sets [30, 42, 59, 110], and Neutrosophic Sets [31, 32, 35, 46, 52–57, 61, 62, 106, 107, 114].

Neutrosophic Sets extend the concept of Fuzzy Sets by incorporating a third dimension—indeterminacy—alongside truth and falsity [104–107]. This approach enables a more nuanced representation of uncertainty and ambiguity. Further advancements have resulted in the development of HyperNeutrosophic Sets and n-SuperHyperNeutrosophic Sets, which are tailored for addressing more intricate and high-dimensional problems [29, 41].

The following subsections present precise definitions and critical attributes of these extended frameworks.

Definition 1.1 (Base Set). A *base set* S is the foundational set from which complex structures such as powersets and hyperstructures are derived. It is formally defined as:

$$S = \{x \mid x \text{ is an element within a specified domain}\}.$$

All elements in constructs like $\mathcal{P}(S)$ or $\mathcal{P}_n(S)$ originate from the elements of S .

Definition 1.2 (Powerset). [37, 94] The *powerset* of a set S , denoted $\mathcal{P}(S)$, is the collection of all possible subsets of S , including both the empty set and S itself. Formally, it is expressed as:

$$\mathcal{P}(S) = \{A \mid A \subseteq S\}.$$

Definition 1.3 (n -th Powerset). (cf. [26, 37, 43, 103, 113])

The n -th powerset of a set H , denoted $P_n(H)$, is defined iteratively, starting with the standard powerset. The recursive construction is given by:

$$P_1(H) = P(H), \quad P_{n+1}(H) = P(P_n(H)), \quad \text{for } n \geq 1.$$

Similarly, the n -th non-empty powerset, denoted $P_n^*(H)$, is defined recursively as:

$$P_1^*(H) = P^*(H), \quad P_{n+1}^*(H) = P^*(P_n^*(H)).$$

Here, $P^*(H)$ represents the powerset of H with the empty set removed.

Definition 1.4 (Neutrosophic Set). [106, 107] Let X be a non-empty set. A *Neutrosophic Set (NS)* A on X is characterized by three membership functions:

$$T_A : X \rightarrow [0, 1], \quad I_A : X \rightarrow [0, 1], \quad F_A : X \rightarrow [0, 1],$$

where for each $x \in X$, the values $T_A(x)$, $I_A(x)$, and $F_A(x)$ represent the degrees of truth, indeterminacy, and falsity, respectively. These values satisfy the following condition:

$$0 \leq T_A(x) + I_A(x) + F_A(x) \leq 3.$$

Definition 1.5 (HyperNeutrosophic Set). (cf. [29, 41, 44, 45, 108]) Let X be a non-empty set. A *HyperNeutrosophic Set (HNS)* \tilde{A} on X is a mapping:

$$\tilde{\mu} : X \rightarrow \mathcal{P}([0, 1]^3),$$

where $\mathcal{P}([0, 1]^3)$ is the family of all non-empty subsets of the unit cube $[0, 1]^3$. For each $x \in X$, $\tilde{\mu}(x) \subseteq [0, 1]^3$ is a set of neutrosophic membership triplets (T, I, F) that satisfy:

$$0 \leq T + I + F \leq 3.$$

Definition 1.6 (n -SuperHyperNeutrosophic Set). (cf. [29, 41, 44, 45]) Let X be a non-empty set. An *n -SuperHyperNeutrosophic Set (n -SHNS)* is a recursive generalization of Neutrosophic Sets and HyperNeutrosophic Sets. It is defined as a mapping:

$$\tilde{A}_n : \mathcal{P}_n(X) \rightarrow \mathcal{P}_n([0, 1]^3),$$

where:

- $\mathcal{P}_1(X) = \mathcal{P}(X)$, the power set of X , and for $k \geq 2$,

$$\mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)),$$

representing the k -th nested family of non-empty subsets of X .

- $\mathcal{P}_n([0, 1]^3)$ is defined similarly for the unit cube $[0, 1]^3$.

For each $A \in \mathcal{P}_n(X)$ and $(T, I, F) \in \tilde{A}_n(A)$, the following condition is satisfied:

$$0 \leq T + I + F \leq 3,$$

where T, I, F represent the degrees of truth, indeterminacy, and falsity for the n -th level subsets of X .

2 Results of This Paper

This section outlines the main results presented in this paper.

2.1 Dymanic HyperNeutrosophic set

A Dynamic Neutrosophic Set incorporates time-dependent truth, indeterminacy, and falsity functions, evolving continuously within a time domain [19, 84, 96, 115–117, 119]. This is extended using Hyperneutrosophic Sets and SuperHyperneutrosophic Sets.

Definition 2.1 (Dynamic Neutrosophic Set). (cf. [116]) Let U be a universal set, and $A \subseteq U$ be a neutrosophic set. A *Dynamic Neutrosophic Set (DNS)* is defined with respect to a time parameter t (where $t \in T, T \subseteq \mathbb{R}^+$) as:

$$D_A^t = \{ \langle x, T_A(t), I_A(t), F_A(t) \rangle \mid x \in U \},$$

where:

- $T_A(t) : U \rightarrow [0, 1]$ is the *truth-membership function* at time t ,
- $I_A(t) : U \rightarrow [0, 1]$ is the *indeterminacy-membership function* at time t ,
- $F_A(t) : U \rightarrow [0, 1]$ is the *falsity-membership function* at time t ,

such that for all $x \in U$,

$$T_A(t) + I_A(t) + F_A(t) \leq 1.$$

The functions $T_A(t), I_A(t), F_A(t)$ are time-dependent and continuous over T . The evolution of the neutrosophic components is represented as:

$$T_A(t), I_A(t), F_A(t) : T \rightarrow [0, 1].$$

Definition 2.2 (Dynamic Hyperneutrosophic Set (DHNS)). Let X be a non-empty set, and let $T \subseteq \mathbb{R}^+$ be a time domain. A *Dynamic Hyperneutrosophic Set (DHNS)* \tilde{D} on X is specified by a mapping:

$$\tilde{D} : X \times T \rightarrow \mathcal{P}([0, 1]^3),$$

such that for each $(x, t) \in X \times T$,

$$\tilde{D}(x, t) \subseteq \{(T, I, F) \in [0, 1]^3 : T + I + F \leq 3\}.$$

Equivalently, for each $t \in T$, the function

$$\tilde{D}_t : X \rightarrow \mathcal{P}([0, 1]^3), \quad \tilde{D}_t(x) := \tilde{D}(x, t),$$

is a Hyperneutrosophic Set in the usual sense, but one that evolves over time t . Each \tilde{D}_t is presumably continuous in t in some sense (optional constraint), reflecting how membership sets might change as t progresses.

Theorem 2.3. *Every Dynamic Neutrosophic Set is a special case of a Dynamic Hyperneutrosophic Set.*

Proof. A *Dynamic Neutrosophic Set (DNS)* \mathcal{D} on U (Definition ??) assigns each (x, t) a single triplet $(T_A(t)(x), I_A(t)(x), F_A(t)(x))$. Let

$$\tilde{D}(x, t) = \{(T_A(t)(x), I_A(t)(x), F_A(t)(x))\}.$$

Thus, each (x, t) maps to a *singleton* in $[0, 1]^3$. We see that $\tilde{D} : X \times T \rightarrow \mathcal{P}([0, 1]^3)$ is a Dynamic Hyperneutrosophic Set. The only difference is that $\tilde{D}(x, t)$ remains single-valued. Hence, every DNS is embedded in DHNS as a degenerate (singleton) membership set for each (x, t) . \square

Theorem 2.4. *Every Hyperneutrosophic Set is a special case of a Dynamic Hyperneutrosophic Set by ignoring time dependence or taking T to be a singleton domain.*

Proof. A *Hyperneutrosophic Set (HNS)* \tilde{A} is a mapping $X \rightarrow \mathcal{P}([0, 1]^3)$. In Definition 2.2, we have $\tilde{D} : X \times T \rightarrow \mathcal{P}([0, 1]^3)$. If we fix $t = t_0$ or let $T = \{t_0\}$ be a single point in time, then

$$\tilde{D}(x, t_0) = \tilde{A}(x).$$

Hence, ignoring or collapsing the time axis recovers the standard hyperneutrosophic membership. Thus, HNS is a sub-case of DHNS where time is trivial or absent. \square

Definition 2.5 (Dynamic n -SuperHyperneutrosophic Set (D- n -SHNS)). Let X be a non-empty set and $T \subseteq \mathbb{R}^+$ be the time domain. Define

$$\mathcal{P}_1(X) = \mathcal{P}(X), \quad \mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)) \quad (k \geq 2).$$

Similarly, let $\mathcal{P}_n([0, 1]^3)$ represent the n -th nested family of non-empty subsets of $[0, 1]^3$. A *Dynamic n -SuperHyperneutrosophic Set (D- n -SHNS)* is a mapping:

$$\tilde{D}_n : \mathcal{P}_n(X) \times T \longrightarrow \mathcal{P}_n([0, 1]^3),$$

such that for each $(A, t) \in \mathcal{P}_n(X) \times T$:

$$\tilde{D}_n(A, t) \subseteq \{(T, I, F) \in [0, 1]^3 : T + I + F \leq 3\}.$$

That is, each n -th level subset A is assigned a *set* of membership triplets (T, I, F) , each lying in $[0, 1]^3$ with $T + I + F \leq 3$, and these sets vary over the time parameter t .

Theorem 2.6. Every Dynamic Hyperneutrosophic Set is a special case of a Dynamic n -SuperHyperneutrosophic Set (D- n -SHNS) for $n = 1$.

Proof. A Dynamic Hyperneutrosophic Set (DHNS) \tilde{D} is a mapping $X \times T \rightarrow \mathcal{P}([0, 1]^3)$. In Definition 2.5, set $n = 1$. Then $\mathcal{P}_1(X) = \mathcal{P}(X)$, and we define

$$\tilde{D}_1(A, t) = \begin{cases} \tilde{D}(x, t), & \text{if } A = \{x\}, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Hence, for singletons $A = \{x\}$, we recover exactly the membership sets from $\tilde{D}(x, t)$. Therefore, a DHNS is embedded in D-1-SHNS as a single-level case. \square

Theorem 2.7. Every (static) n -SuperHyperneutrosophic Set is a special case of a Dynamic n -SuperHyperneutrosophic Set by taking T to be a single point or ignoring time.

Proof. An n -SuperHyperneutrosophic Set (SHNS) \tilde{A}_n is a mapping $\mathcal{P}_n(X) \rightarrow \mathcal{P}_n([0, 1]^3)$. In Definition 2.5, we have $\tilde{D}_n : \mathcal{P}_n(X) \times T \rightarrow \mathcal{P}_n([0, 1]^3)$. If we let $T = \{t_0\}$ be a single point in time (or otherwise disregard time), we can define

$$\tilde{D}_n(A, t_0) = \tilde{A}_n(A).$$

Then ignoring the time dimension yields the standard n -SuperHyperneutrosophic membership sets. Thus, every n -SHNS is included in D- n -SHNS by collapsing T to a singleton or skipping time dependence. \square

2.2 Hyper Quadripartitioned Neutrosophic set

A Quadripartitioned Neutrosophic Set assigns four membership values [16, 17, 23, 49, 69, 70, 86, 88, 99–101, 104]. This is extended using Hyperneutrosophic Sets and SuperHyperneutrosophic Sets.

Definition 2.8 (Quadripartitioned Neutrosophic Set (QNS)). (cf. [16, 23, 69]) Let X be a universe of discourse. A *Quadripartitioned Neutrosophic Set (QNS)* on X is given by

$$QNS = \{\langle x, T(x), C(x), U(x), F(x) \rangle \mid x \in X\},$$

where each of $T(x), C(x), U(x), F(x)$ lies in $[0, 1]$, satisfying

$$0 \leq T(x) + C(x) + U(x) + F(x) \leq 4.$$

Definition 2.9 (Hyper Quadripartitioned Neutrosophic Set (HQNS)). Let X be a non-empty set, and consider the family $\mathcal{P}([0, 1]^4)$ of all non-empty subsets of the unit 4-cube $[0, 1]^4$. A *Hyper Quadripartitioned Neutrosophic Set (HQNS)* \tilde{Q} on X is a mapping

$$\tilde{Q} : X \longrightarrow \mathcal{P}([0, 1]^4),$$

such that for each $x \in X$,

$$\tilde{Q}(x) \subseteq \{(T, C, U, F) \in [0, 1]^4 : T + C + U + F \leq 4\}.$$

That is, each point $x \in X$ is assigned a *set* of quadripartitioned membership quadruples (T, C, U, F) , each lying in $[0, 1]^4$ with $T + C + U + F \leq 4$.

Theorem 2.10. *Every Quadripartitioned Neutrosophic Set is a special case of a Hyper Quadripartitioned Neutrosophic Set.*

Proof. A Quadripartitioned Neutrosophic Set (QNS) Q assigns each $x \in X$ exactly one quadruple

$$(T(x), C(x), U(x), F(x)) \in [0, 1]^4$$

with $T + C + U + F \leq 4$. In Definition 2.9, we let

$$\tilde{Q}(x) = \{(T(x), C(x), U(x), F(x))\},$$

a singleton subset of $[0, 1]^4$. The same constraint $T + C + U + F \leq 4$ persists, so each single-valued QNS is embedded in the HQNS framework as a degenerate (singleton) membership set for each x . \square

Theorem 2.11. *Every HyperNeutrosophic Set is a special case of a Hyper Quadripartitioned Neutrosophic Set by treating the four components as, for instance, $(T, I, 0, F)$ (or by ignoring C and U).*

Proof. A HyperNeutrosophic Set (HNS) \tilde{A} maps each $x \in X$ to a subset of $[0, 1]^3$ with $T + I + F \leq 3$. In Definition 2.9, each membership is a subset of $[0, 1]^4$ with $T + C + U + F \leq 4$. If we fix $C = 0$ and $U = I$ (or $U = 0$) so that effectively (T, C, U, F) becomes $(T, 0, I, F)$, and require $T + 0 + I + F = T + I + F \leq 3$, we see that ignoring or collapsing the extra dimension recovers an HNS membership set. Specifically, define

$$\tilde{Q}(x) = \{(T, I, 0, F) : (T, I, F) \in \tilde{A}(x)\}.$$

Hence, ignoring two of the four components (or setting them to zero or merging them) yields the usual 3-component hyperneutrosophic membership. Therefore, an HNS is subsumed under HQNS by discarding or collapsing extra components to zero. \square

Definition 2.12 (n -SuperHyper Quadripartitioned Neutrosophic Set (n -SHQNS)). Let X be a non-empty set. Define:

$$\mathcal{P}_1(X) = \mathcal{P}(X), \quad \mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)) \quad (k \geq 2).$$

Similarly, let $\mathcal{P}_n([0, 1]^4)$ denote the n -th nested family of non-empty subsets of $[0, 1]^4$. A n -SuperHyper Quadripartitioned Neutrosophic Set (n -SHQNS) is a mapping

$$\tilde{Q}_n : \mathcal{P}_n(X) \longrightarrow \mathcal{P}_n([0, 1]^4),$$

such that for each $A \in \mathcal{P}_n(X)$,

$$\tilde{Q}_n(A) \subseteq \{(T, C, U, F) \in [0, 1]^4 : T + C + U + F \leq 4\}.$$

Hence, each n -th level subset $A \subseteq X$ is assigned a set of four-part membership quadruples (T, C, U, F) in $[0, 1]^4$ with $T + C + U + F \leq 4$.

Theorem 2.13. *Every Hyper Quadripartitioned Neutrosophic Set is a special case of an n -SuperHyper Quadripartitioned Neutrosophic Set (n -SHQNS) for $n = 1$.*

Proof. A Hyper Quadripartitioned Neutrosophic Set (HQNS) \tilde{Q} is a mapping $X \rightarrow \mathcal{P}([0, 1]^4)$ with $T + C + U + F \leq 4$. In Definition 2.12, set $n = 1$, so

$$\tilde{Q}_1 : \mathcal{P}_1(X) = \mathcal{P}(X) \rightarrow \mathcal{P}_1([0, 1]^4) = \mathcal{P}([0, 1]^4).$$

Define

$$\tilde{Q}_1(\{x\}) := \tilde{Q}(x), \quad \tilde{Q}_1(A) = \emptyset \quad \text{for } A \neq \{x\}.$$

Hence, for singletons $A = \{x\} \subseteq X$, we recover precisely $\tilde{Q}(x)$. The constraint $T + C + U + F \leq 4$ remains the same, embedded in $\tilde{Q}_1(A)$. Therefore, any HQNS is realized as a degenerate single-level mapping in an n -SHQNS with $n = 1$. \square

Theorem 2.14. *Every n -SuperHyperNeutrosophic Set is a special case of an n -SuperHyper Quadripartitioned Neutrosophic Set by ignoring or collapsing the fourth component (e.g., $C = 0$ or $U = I$).*

Proof. An n -SuperHyperNeutrosophic Set (SHNS) \tilde{A}_n maps each $A \in \mathcal{P}_n(X)$ to a set of (T, I, F) in $[0, 1]^3$ with $T + I + F \leq 3$. In Definition 2.12, each membership is in $[0, 1]^4$ with $T + C + U + F \leq 4$. To retrieve an SHNS, fix or zero out some components. For instance, set $C = 0$, $U = I$, and require $T + I + F = T + (U) + F \leq 3$. Concretely, define

$$\tilde{Q}_n(A) = \{(T, 0, I, F) : (T, I, F) \in \tilde{A}_n(A)\}.$$

Hence, ignoring the extra dimension(s) reverts membership to $(T, I, F) \in [0, 1]^3$. Therefore, an n -SHNS is a special case of n -SHQNS with additional constraints or identification of dimensions. \square

2.3 Hyper Pentapartitioned Neutrosophic set

A Pentapartitioned Neutrosophic Set assigns five membership values [5, 13, 15, 20–22, 28, 33, 36, 40, 58, 76, 85, 87]. This is extended using Hyperneutrosophic Sets and SuperHyperneutrosophic Sets.

Definition 2.15 (Pentapartitioned Neutrosophic Set (PNS)). (cf. [13, 76, 85]) Let X be a universe of discourse. A Pentapartitioned Neutrosophic Set (PNS) on X is given by

$$PNS = \{(x, T(x), C(x), R(x), U(x), F(x)) \mid x \in X\},$$

where each of $T(x), C(x), R(x), U(x), F(x) \in [0, 1]$, satisfying

$$0 \leq T(x) + C(x) + R(x) + U(x) + F(x) \leq 5.$$

Definition 2.16 (Hyper Pentapartitioned Neutrosophic Set (HPNS)). Let X be a non-empty set, and let $\mathcal{P}([0, 1]^5)$ denote the family of all non-empty subsets of the unit 5-cube $[0, 1]^5$. A Hyper Pentapartitioned Neutrosophic Set (HPNS) \tilde{P} on X is a mapping

$$\tilde{P} : X \rightarrow \mathcal{P}([0, 1]^5),$$

such that for each $x \in X$,

$$\tilde{P}(x) \subseteq \{(T, C, R, U, F) \in [0, 1]^5 : T + C + R + U + F \leq 5\}.$$

Hence, each $x \in X$ is assigned a set of pentapartitioned membership quintuples (T, C, R, U, F) , where $T + C + R + U + F \leq 5$.

Theorem 2.17. *Every Pentapartitioned Neutrosophic Set is a special case of a Hyper Pentapartitioned Neutrosophic Set.*

Proof. A Pentapartitioned Neutrosophic Set (PNS) assigns each $x \in X$ a unique quintuple

$$(T(x), C(x), R(x), U(x), F(x)) \in [0, 1]^5$$

satisfying $T + C + R + U + F \leq 5$. In Definition 2.16, we map each x to a set of such quintuples. Define:

$$\tilde{P}(x) = \{(T(x), C(x), R(x), U(x), F(x))\},$$

i.e. a singleton set. The same constraint $T + C + R + U + F \leq 5$ persists. Consequently, each PNS is embedded in HPNS as a degenerate (singleton) membership set for each x . \square

Theorem 2.18. *Every HyperNeutrosophic Set is a special case of a Hyper Pentapartitioned Neutrosophic Set by reducing two membership components to zero (e.g., $C = R = 0$).*

Proof. A *HyperNeutrosophic Set (HNS)* \tilde{A} maps each $x \in X$ to a subset of $[0, 1]^3$ (triplets (T, I, F) with $T + I + F \leq 3$). In Definition 2.16, each membership is a subset of $[0, 1]^5$ with $T + C + R + U + F \leq 5$. If we identify (T, I, F) in $[0, 1]^3$ with $(T, 0, 0, I, F)$ in $[0, 1]^5$, we require $T + 0 + 0 + I + F = T + I + F \leq 3$. We can embed an HNS as:

$$\tilde{P}(x) = \{(T, 0, 0, I, F) : (T, I, F) \in \tilde{A}(x)\}.$$

Hence, ignoring or zeroing out two membership parts ($C = 0, R = 0$) recovers a 3-part hyperneutrosophic membership. Therefore, an HNS is included in HPNS by discarding extra membership dimensions. \square

Definition 2.19 (*n-SuperHyper Pentapartitioned Neutrosophic Set (n-SHPNS)*). Let X be a non-empty set. Define recursively:

$$\mathcal{P}_1(X) = \mathcal{P}(X), \quad \mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)) \quad \text{for } k \geq 2.$$

Similarly, let $\mathcal{P}_n([0, 1]^5)$ denote the n -th nested family of non-empty subsets of $[0, 1]^5$. A *n-SuperHyper Pentapartitioned Neutrosophic Set (n-SHPNS)* is a mapping

$$\tilde{P}_n : \mathcal{P}_n(X) \longrightarrow \mathcal{P}_n([0, 1]^5),$$

such that for each $A \in \mathcal{P}_n(X)$,

$$\tilde{P}_n(A) \subseteq \{(T, C, R, U, F) \in [0, 1]^5 : T + C + R + U + F \leq 5\}.$$

Hence, each n -th level subset A is assigned a *set* of pentapartitioned membership quintuples $(T, C, R, U, F) \in [0, 1]^5$, with $T + C + R + U + F \leq 5$.

Theorem 2.20. *Every Hyper Pentapartitioned Neutrosophic Set is a special case of an n-SuperHyper Pentapartitioned Neutrosophic Set (n-SHPNS) for $n = 1$.*

Proof. A *Hyper Pentapartitioned Neutrosophic Set (HPNS)* \tilde{P} is a mapping $X \rightarrow \mathcal{P}([0, 1]^5)$ with $T + C + R + U + F \leq 5$. In Definition 2.19, set $n = 1$. Then

$$\tilde{P}_1 : \mathcal{P}_1(X) = \mathcal{P}(X) \rightarrow \mathcal{P}_1([0, 1]^5) = \mathcal{P}([0, 1]^5).$$

Define

$$\tilde{P}_1(\{x\}) := \tilde{P}(x), \quad \tilde{P}_1(A) = \emptyset \quad (\text{for } A \neq \{x\}).$$

Hence, for singletons $A = \{x\} \subseteq X$, we recover exactly $\tilde{P}(x)$. The constraint $T + C + R + U + F \leq 5$ is maintained. Therefore, any HPNS is subsumed in n-SHPNS with $n = 1$. \square

Theorem 2.21. *Every n-SuperHyperneutrosophic Set is a special case of an n-SuperHyper Pentapartitioned Neutrosophic Set by reducing two membership components to zero (e.g., $C = R = 0$).*

Proof. An *n-SuperHyperneutrosophic Set (SHNS)* \tilde{A}_n maps each $A \in \mathcal{P}_n(X)$ to a subset of $[0, 1]^3$ (triplets (T, I, F)). Compare with Definition 2.19, where $\tilde{P}_n(A)$ is a subset of $[0, 1]^5$ with $T + C + R + U + F \leq 5$. To recover an SHNS from n-SHPNS, we set $C = R = 0$ and identify $U = I$. Hence $(T, C, R, U, F) = (T, 0, 0, I, F)$ with $T + I + F \leq 3$. Formally, define

$$\tilde{P}_n(A) = \{(T, 0, 0, I, F) : (T, I, F) \in \tilde{A}_n(A)\}.$$

Hence, ignoring or zeroing out two membership parts yields a standard 3-part membership in $[0, 1]^3$. Thus, each SHNS is included in an n-SHPNS by discarding extra membership dimensions. \square

2.4 Hyper Heptapartitioned Neutrosophic Set

A Heptapartitioned Neutrosophic Set assigns seven membership [14, 44, 80–82]. This is extended using Hyperneutrosophic Sets and SuperHyperneutrosophic Sets.

Definition 2.22 (Heptapartitioned Neutrosophic Set). [14, 81] A Heptapartitioned Neutrosophic Set (HNS) on a universe X is defined as:

$$HNS = \{ \langle x, T(x), C(x), R(x), U(x), F(x), G(x), L(x) \rangle \mid x \in X \},$$

where $T(x), C(x), R(x), U(x), F(x), G(x), L(x) \in [0, 1]$, and

$$0 \leq T(x) + C(x) + R(x) + U(x) + F(x) + G(x) + L(x) \leq 7.$$

Definition 2.23 (Hyper Heptapartitioned Neutrosophic Set (HHNS)). Let X be a non-empty set, and let $\mathcal{P}([0, 1]^7)$ denote the family of all non-empty subsets of the 7-cube $[0, 1]^7$. A Hyper Heptapartitioned Neutrosophic Set (HHNS) \tilde{H} on X is a mapping

$$\tilde{H} : X \longrightarrow \mathcal{P}([0, 1]^7),$$

such that for each $x \in X$,

$$\tilde{H}(x) \subseteq \{ (T, C, R, U, F, G, L) \in [0, 1]^7 : T + C + R + U + F + G + L \leq 7 \}.$$

Hence, every point $x \in X$ is assigned a set of heptapartitioned membership 7-tuples (T, C, R, U, F, G, L) , each lying in $[0, 1]^7$ with $T + C + R + U + F + G + L \leq 7$.

Theorem 2.24. Every Heptapartitioned Neutrosophic Set is a special case of a Hyper Heptapartitioned Neutrosophic Set.

Proof. A Heptapartitioned Neutrosophic Set (HptNS) H assigns each $x \in X$ exactly one 7-tuple $(T, C, R, U, F, G, L) \in [0, 1]^7$ with $T + C + R + U + F + G + L \leq 7$. To embed this in Definition 2.23, define

$$\tilde{H}(x) = \left\{ (T(x), C(x), R(x), U(x), F(x), G(x), L(x)) \right\},$$

i.e. a singleton set in $[0, 1]^7$. Since the same $T + C + R + U + F + G + L \leq 7$ constraint remains, each single-valued HptNS is embedded in HHNS as a degenerate membership set (a singleton). \square

Theorem 2.25. Every HyperNeutrosophic Set is a special case of a Hyper Heptapartitioned Neutrosophic Set by ignoring four of the membership components.

Proof. A HyperNeutrosophic Set (HNS) \tilde{A} maps each $x \in X$ to a subset of $[0, 1]^3$ (triplets (T, I, F)). In Definition 2.23, each membership is a subset of $[0, 1]^7$ with $T + C + R + U + F + G + L \leq 7$. If we fix four components to zero (e.g., $C = R = G = L = 0$) and rename $U = I$, then (T, I, F) in $[0, 1]^3$ becomes $(T, 0, 0, I, F, 0, 0)$ in $[0, 1]^7$, needing $T + 0 + 0 + I + F + 0 + 0 = T + I + F \leq 3$. Formally:

$$\tilde{H}(x) = \left\{ (T, 0, 0, I, F, 0, 0) : (T, I, F) \in \tilde{A}(x) \right\}.$$

Hence, ignoring the extra membership components merges an HNS into an HHNS. Therefore, an HNS is a special case of HHNS by discarding (or zeroing out) the four additional partitions. \square

Definition 2.26 (n -SuperHyper Heptapartitioned Neutrosophic Set (n -SHHNS)). Let X be a non-empty set. Define recursively:

$$\mathcal{P}_1(X) = \mathcal{P}(X), \quad \mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)) \quad (k \geq 2).$$

Similarly, let $\mathcal{P}_n([0, 1]^7)$ denote the n -th nested family of non-empty subsets of the 7-cube $[0, 1]^7$. A n -SuperHyper Heptapartitioned Neutrosophic Set (n -SHHNS) is a mapping

$$\tilde{H}_n : \mathcal{P}_n(X) \rightarrow \mathcal{P}_n([0, 1]^7),$$

such that for each $A \in \mathcal{P}_n(X)$,

$$\tilde{H}_n(A) \subseteq \{ (T, C, R, U, F, G, L) \in [0, 1]^7 : T + C + R + U + F + G + L \leq 7 \}.$$

Hence, every n -th level subset A is assigned a set of membership 7-tuples $(T, C, R, U, F, G, L) \in [0, 1]^7$ with $T + C + R + U + F + G + L \leq 7$.

Theorem 2.27. Every Hyper Heptapartitioned Neutrosophic Set is a special case of an n -SuperHyper Heptapartitioned Neutrosophic Set for $n = 1$.

Proof. A Hyper Heptapartitioned Neutrosophic Set (HHNS) \tilde{H} is a mapping $X \rightarrow \mathcal{P}([0, 1]^7)$ with $T + C + R + U + F + G + L \leq 7$. In Definition 2.26, set $n = 1$, so

$$\tilde{H}_1 : \mathcal{P}_1(X) = \mathcal{P}(X) \rightarrow \mathcal{P}_1([0, 1]^7) = \mathcal{P}([0, 1]^7).$$

Define

$$\tilde{H}_1(\{x\}) := \tilde{H}(x), \quad \tilde{H}_1(A) = \emptyset \quad (\text{for } A \neq \{x\}).$$

Hence, for singletons $A = \{x\} \subseteq X$, we recover precisely the membership sets $\tilde{H}(x)$. Thus, every HHNS is embedded in an n -SHHNS with $n = 1$. \square

Theorem 2.28. Every n -SuperHyperneutrosophic Set is a special case of an n -SuperHyper Heptapartitioned Neutrosophic Set by ignoring four membership components (e.g. $C = R = G = L = 0$).

Proof. An n -SuperHyperneutrosophic Set (SHNS) \tilde{A}_n maps each $A \in \mathcal{P}_n(X)$ to subsets of $[0, 1]^3$ with (T, I, F) membership. In Definition 2.26, we assign subsets of $[0, 1]^7$ with (T, C, R, U, F, G, L) , each summing to at most 7. If we require $C = R = G = L = 0$ and rename $U = I$, then (T, I, F) in $[0, 1]^3$ is identified with $(T, 0, 0, I, F, 0, 0)$ in $[0, 1]^7$. So define

$$\tilde{H}_n(A) = \left\{ (T, 0, 0, I, F, 0, 0) : (T, I, F) \in \tilde{A}_n(A) \right\}.$$

Hence, ignoring four extra membership dimensions reverts us to $(T, I, F) \in [0, 1]^3$. Therefore, an SHNS is included in n -SHHNS by collapsing the four additional components. \square

2.5 m-Polar Hyperneutrosophic Set

An m -Polar Hyperneutrosophic Set extends the conventional neutrosophic framework by assigning m distinct truth, indeterminacy, and falsity triplets to each element in a given universe. This model provides a comprehensive structure for handling complex and multidimensional uncertainties [67, 75, 78, 83, 95, 98, 99, 102]. Related concepts include the bipolar neutrosophic set [1, 24, 77, 118], bipolar fuzzy set [3, 4, 66], tripolar fuzzy set [89–91], and m -polar fuzzy set [18, 63, 92, 93], which address specific dimensions of uncertainty and vagueness.

The m -Polar Neutrosophic Set is further extended using the frameworks of Hyperneutrosophic Sets and SuperHyperneutrosophic Sets, allowing for even more flexible and detailed representations of complex systems.

Definition 2.29 (m -Polar Neutrosophic Set). (cf. [83, 95, 98, 99, 102]) Let X be a universe of discourse and $m \geq 1$ represent the number of poles or criteria. An m -polar neutrosophic set A is defined as:

$$A = \left\{ \langle x, (T_A^{(k)}(x), I_A^{(k)}(x), F_A^{(k)}(x)) \rangle \mid x \in X, k = 1, 2, \dots, m \right\}, \text{ AbdelBasset2019CosineSM},$$

where:

- $T_A^{(k)}(x), I_A^{(k)}(x), F_A^{(k)}(x) \in [0, 1]$,
- $T_A^{(k)}(x) + I_A^{(k)}(x) + F_A^{(k)}(x) \leq 3, \forall x \in X, k = 1, 2, \dots, m$.

Definition 2.30 (m -Polar Hyperneutrosophic Set (m -HNS)). Let X be a non-empty set, and let $m \geq 1$. Consider the family $\mathcal{P}([0, 1]^3)$ of all non-empty subsets of the unit cube $[0, 1]^3$. An m -Polar Hyperneutrosophic Set \tilde{M} on X is defined by a mapping

$$\tilde{M} : X \times \{1, 2, \dots, m\} \rightarrow \mathcal{P}([0, 1]^3),$$

such that for each (x, k) :

$$\tilde{M}(x, k) \subseteq \{(T, I, F) \in [0, 1]^3 : T + I + F \leq 3\}.$$

Hence, each element $x \in X$ and each pole k is assigned a set of possible triplets (T, I, F) .

Theorem 2.31. *Every m -Polar Neutrosophic Set is a special case of an m -Polar Hyperneutrosophic Set.*

Proof. An m -Polar Neutrosophic Set (mPNS) A assigns each (x, k) exactly one triplet $(T_A^{(k)}(x), I_A^{(k)}(x), F_A^{(k)}(x))$ with $T + I + F \leq 3$. In Definition 2.30, we let each (x, k) map to a set in $[0, 1]^3$. Define

$$\tilde{M}(x, k) = \left\{ (T_A^{(k)}(x), I_A^{(k)}(x), F_A^{(k)}(x)) \right\},$$

a singleton. The usual constraint $T + I + F \leq 3$ holds. Hence, every mPNS is embedded in an m -Polar Hyperneutrosophic Set as a degenerate case (singleton membership for each (x, k)). \square

Theorem 2.32. *Every HyperNeutrosophic Set is a special case of an m -Polar Hyperneutrosophic Set by setting $m = 1$ (only one pole).*

Proof. A HyperNeutrosophic Set (HNS) \tilde{A} is a mapping $X \rightarrow \mathcal{P}([0, 1]^3)$. In Definition 2.30, we have $\tilde{M} : X \times \{1, 2, \dots, m\} \rightarrow \mathcal{P}([0, 1]^3)$. If $m = 1$, then for each x we define

$$\tilde{M}(x, 1) = \tilde{A}(x).$$

Hence, ignoring the multiple poles (just $k = 1$) yields exactly a HyperNeutrosophic Set. Therefore, an HNS is a sub-case of an m -Polar Hyperneutrosophic Set with $m = 1$. \square

Definition 2.33 (m -Polar n -SuperHyperneutrosophic Set (m-SHNS)). Let X be a non-empty set, $m \geq 1$ be the number of poles, and define recursively:

$$\mathcal{P}_1(X) = \mathcal{P}(X), \quad \mathcal{P}_k(X) = \mathcal{P}(\mathcal{P}_{k-1}(X)) \quad (k \geq 2).$$

Similarly, consider $\mathcal{P}_n([0, 1]^3)$ for n -nested subsets of $[0, 1]^3$. An m -Polar n -SuperHyperneutrosophic Set \tilde{M}_n is given by a mapping

$$\tilde{M}_n : \mathcal{P}_n(X) \times \{1, 2, \dots, m\} \longrightarrow \mathcal{P}_n([0, 1]^3),$$

such that for each $(A, k) \in \mathcal{P}_n(X) \times \{1, \dots, m\}$:

$$\tilde{M}_n(A, k) \subseteq \{(T, I, F) \in [0, 1]^3 : T + I + F \leq 3\}.$$

Hence, each n -th level subset A and each pole k is assigned a set of membership triplets $(T, I, F) \in [0, 1]^3$.

Theorem 2.34. *Every m -Polar Hyperneutrosophic Set is a special case of an m -Polar n -SuperHyperneutrosophic Set for $n = 1$.*

Proof. An m -Polar Hyperneutrosophic Set \tilde{M} (Definition 2.30) is a mapping:

$$X \times \{1, \dots, m\} \rightarrow \mathcal{P}([0, 1]^3).$$

In Definition 2.33, for $n = 1$ we have $\tilde{M}_1 : \mathcal{P}_1(X) \times \{1, \dots, m\} \rightarrow \mathcal{P}_1([0, 1]^3) = \mathcal{P}([0, 1]^3)$. We can embed \tilde{M} by defining:

$$\tilde{M}_1(\{x\}, k) := \tilde{M}(x, k), \quad \tilde{M}_1(A, k) = \emptyset \quad \text{if } A \neq \{x\}.$$

Hence, for singletons $\{x\} \subset X$, we recover exactly the membership sets from $\tilde{M}(x, k)$. The constraint $T + I + F \leq 3$ remains identical. Therefore, \tilde{M} is included in \tilde{M}_1 , which is an m -Polar 1-SuperHyperneutrosophic Set. \square

Theorem 2.35. *Every n -SuperHyperneutrosophic Set is a special case of an m -Polar n -SuperHyperneutrosophic Set by letting $m = 1$ (only one pole).*

Proof. An n -SuperHyperneutrosophic Set (SHNS) \tilde{A}_n is a mapping $\mathcal{P}_n(X) \rightarrow \mathcal{P}_n([0, 1]^3)$. In Definition 2.33, we have $\tilde{M}_n : \mathcal{P}_n(X) \times \{1, \dots, m\} \rightarrow \mathcal{P}_n([0, 1]^3)$. If $m = 1$, we define

$$\tilde{M}_n(A, 1) = \tilde{A}_n(A), \quad \tilde{M}_n(A, k) = \emptyset \quad \text{for } k \neq 1.$$

Hence, ignoring multiple poles (just $k = 1$) we recover the usual n -SuperHyperneutrosophic mapping from $\mathcal{P}_n(X)$ to $\mathcal{P}_n([0, 1]^3)$. Therefore, an SHNS is embedded in m -SHNS with $m = 1$. \square

Funding

This study did not receive any financial or external support from organizations or individuals.

Acknowledgments

We extend our sincere gratitude to everyone who provided insights, inspiration, and assistance throughout this research. We particularly thank our readers for their interest and acknowledge the authors of the cited works for laying the foundation that made our study possible. We also appreciate the support from individuals and institutions that provided the resources and infrastructure needed to produce and share this paper. Finally, we are grateful to all those who supported us in various ways during this project.

Data Availability

This research is purely theoretical, involving no data collection or analysis. We encourage future researchers to pursue empirical investigations to further develop and validate the concepts introduced here.

Ethical Approval

As this research is entirely theoretical in nature and does not involve human participants or animal subjects, no ethical approval is required.

Conflicts of Interest

The authors confirm that there are no conflicts of interest related to the research or its publication.

Disclaimer

This work presents theoretical concepts that have not yet undergone practical testing or validation. Future researchers are encouraged to apply and assess these ideas in empirical contexts. While every effort has been made to ensure accuracy and appropriate referencing, unintentional errors or omissions may still exist. Readers are advised to verify referenced materials on their own. The views and conclusions expressed here are the authors' own and do not necessarily reflect those of their affiliated organizations.

References

- [1] Mohamed Abdel-Basset, Mai Mohamed, Mohamed Elhoseny, Le Hoang Son, Francisco Chiclana, and Abdel Nasser H. Zaied. Cosine similarity measures of bipolar neutrosophic set for diagnosis of bipolar disorder diseases. *Artificial intelligence in medicine*, 101:101735, 2019.
- [2] Muhammad Akram, A Nagoor Gani, and A Borumand Saeid. Vague hypergraphs. *Journal of Intelligent & Fuzzy Systems*, 26(2):647–653, 2014.
- [3] Muhammad Akram, Musavarah Sarwar, Wieslaw A Dudek, Muhammad Akram, Musavarah Sarwar, and Wieslaw A Dudek. Bipolar neutrosophic graph structures. *Graphs for the Analysis of Bipolar Fuzzy Information*, pages 393–446, 2021.
- [4] Muhammad Akram, Shumaiza, and José Carlos Rodríguez Alcantud. Multi-criteria decision making methods with bipolar fuzzy sets. *Forum for Interdisciplinary Mathematics*, 2023.
- [5] Manal Al-Labadi, Shuker Khalil, VR Radhika, K Mohana, et al. Pentapartitioned neutrosophic vague soft sets and its applications. *International Journal of Neutrosophic Science*, (2):64–4, 2025.
- [6] Krassimir Atanassov. Intuitionistic fuzzy sets. *International journal bioautomation*, 20:1, 2016.
- [7] Krassimir Atanassov and George Gargov. Elements of intuitionistic fuzzy logic. part i. *Fuzzy sets and systems*, 95(1):39–52, 1998.
- [8] Krassimir T Atanassov. *On intuitionistic fuzzy sets theory*, volume 283. Springer, 2012.
- [9] Krassimir T Atanassov. Circular intuitionistic fuzzy sets. *Journal of Intelligent & Fuzzy Systems*, 39(5):5981–5986, 2020.
- [10] Krassimir T Atanassov and Krassimir T Atanassov. *Intuitionistic fuzzy sets*. Springer, 1999.
- [11] Krassimir T Atanassov and G Gargov. *Intuitionistic fuzzy logics*. Springer, 2017.
- [12] Vivek Badhe, RS Thakur, and GS Thakur. Vague set theory for profit pattern and decision making in uncertain data. *International journal of advanced computer science and applications*, 6(6):58–64, 2015.

-
- [13] Pranab Biswas, Surapati Pramanik, and Bibhas Chandra Giri. Single valued bipolar pentapartitioned neutrosophic set and its application in madm strategy. 2022.
 - [14] S Broumi and Tomasz Witczak. Heptapartitioned neutrosophic soft set. *International Journal of Neutrosophic Science*, 18(4):270–290, 2022.
 - [15] Said Broumi, D Ajay, P Chellamani, Lathamaheswari Malayalan, Mohamed Talea, Assia Bakali, Philippe Schweizer, and Saeid Jafari. Interval valued pentapartitioned neutrosophic graphs with an application to mcdm. *Operational Research in Engineering Sciences: Theory and Applications*, 5(3):68–91, 2022.
 - [16] Rajashi Chatterjee, Pinaki Majumdar, and Syamal Kumar Samanta. A multi-criteria group decision making algorithm with quadri-partitioned neutrosophic weighted aggregation operators using quadripartitioned neutrosophic numbers in ipqsvnss environment. *Soft Computing*, 24(12):8857–8880, 2020.
 - [17] Tanmoy Chatterjee and Surapati Pramanik. Triangular fuzzy quadripartitioned neutrosophic set and its properties. *Neutrosophic Sets and Systems*, 75:15–28, 2025.
 - [18] Juanjuan Chen, Shenggang Li, Shengquan Ma, and Xueping Wang. m-polar fuzzy sets: an extension of bipolar fuzzy sets. *The scientific world journal*, 2014(1):416530, 2014.
 - [19] Wen-Hua Cui and Jun Ye. Logarithmic similarity measure of dynamic neutrosophic cubic sets and its application in medical diagnosis. *Computers in Industry*, 111:198–206, 2019.
 - [20] Rakhal Das and Suman Das. Pentapartitioned neutrosophic subtraction algebra. *Neutrosophic Sets and Systems*, 68:89–98, 2024.
 - [21] Suman Das, Rakhal Das, and Binod Chandra Tripathy. Topology on rough pentapartitioned neutrosophic set. *Iraqi Journal of Science*, 2022.
 - [22] Suman Das, Prasanna Poojary, Vadiraja Bhatta GR, and Sunny Kumar Sharma. Single-valued pentapartitioned neutrosophic bi-topological spaces. *International Journal of Neutrosophic Science (IJNS)*, 23(4), 2024.
 - [23] Somen Debnath. Fuzzy quadripartitioned neutrosophic soft matrix theory and its decision-making approach. *Journal of Computational and Cognitive Engineering*, 1(2):88–93, 2022.
 - [24] Irfan Deli, Mumtaz Ali, and Florentin Smarandache. Bipolar neutrosophic sets and their application based on multi-criteria decision making problems. *2015 International Conference on Advanced Mechatronic Systems (ICAMechS)*, pages 249–254, 2015.
 - [25] Ronald C. Freiwald. An introduction to set theory and topology. 2014.
 - [26] Takaaki Fujita. A concise review on various concepts of superhyperstructures.
 - [27] Takaaki Fujita. Natural n-superhyper plithogenic language.
 - [28] Takaaki Fujita. Review of plithogenic directed, mixed, bidirected, and pangene offgraph. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 120.
 - [29] Takaaki Fujita. Exploring concepts of hyperfuzzy, hyperneutrosophic, and hyperplithogenic sets. 2024. DOI: 10.13140/RG.2.2.12216.87045.
 - [30] Takaaki Fujita. Note for hypersoft filter and fuzzy hypersoft filter. *Multicriteria Algorithms With Applications*, 5:32–51, 2024.
 - [31] Takaaki Fujita. Note for neutrosophic incidence and threshold graph. *SciNexus*, 1:97–125, 2024.
 - [32] Takaaki Fujita. Reconsideration of neutrosophic social science and neutrosophic phenomenology with non-classical logic. Technical report, Center for Open Science, 2024.
 - [33] Takaaki Fujita. A review of fuzzy and neutrosophic offsets: Connections to some set concepts and normalization function. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 74, 2024.
 - [34] Takaaki Fujita. Short communication of big plithogenic science and deep plithogenic science. 2024. preprint (researchgate).
 - [35] Takaaki Fujita. Short note of bunch graph in fuzzy, neutrosophic, and plithogenic graphs. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 294, 2024.
 - [36] Takaaki Fujita. Some graph classes for turiyam neutrosophic directed graphs and pentapartitioned neutrosophic directed graphs. 2024.
 - [37] Takaaki Fujita. Superhypergraph neural networks and plithogenic graph neural networks: Theoretical foundations. *arXiv preprint arXiv:2412.01176*, 2024.
 - [38] Takaaki Fujita. Survey of intersection graphs, fuzzy graphs and neutrosophic graphs. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 114, 2024.
 - [39] Takaaki Fujita. Survey of planar and outerplanar graphs in fuzzy and neutrosophic graphs. *ResearchGate*, July 2024.
 - [40] Takaaki Fujita. Uncertain labeling graphs and uncertain graph classes (with survey for various uncertain sets). July 2024.
 - [41] Takaaki Fujita. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*. Biblio Publishing, 2025.
 - [42] Takaaki Fujita. A comprehensive discussion on fuzzy hypersoft expert, superhypersoft, and indeterminsoft graphs. *Neutrosophic Sets and Systems*, 77:241–263, 2025.
 - [43] Takaaki Fujita. Exploration of graph classes and concepts for superhypergraphs and n-th power mathematical structures. 2025.
 - [44] Takaaki Fujita. Exploring concepts of hyperfuzzy, hyperneutrosophic, and hyperplithogenic sets ii. *ResearchGate*, 2025.
 - [45] Takaaki Fujita. Hyperfuzzy hyperrough set, hyperneutrosophic hyperrough set, and hypersoft hyperrough set. Preprint, 2025.
 - [46] Takaaki Fujita. Short note of even-hole-graph for uncertain graph. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 351, 2025.
 - [47] Takaaki Fujita. Short note of extended hyperplithogenic sets, 2025. Preprint.

-
- [48] Takaaki Fujita. A short note on the basic graph construction algorithm for plithogenic graphs. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 274, 2025.
 - [49] Takaaki Fujita. Short survey on the hierarchical uncertainty of fuzzy, neutrosophic, and plithogenic sets, 2025. Preprint.
 - [50] Takaaki Fujita. Some type of hyperneutrosophic set: Bipolar, pythagorean, double-valued, interval-valued set, 2025. Preprint.
 - [51] Takaaki Fujita. Some types of hyperneutrosophic set (2): Complex, single-valued triangular, fermatean, and linguistic sets. *Preprint*, 2025.
 - [52] Takaaki Fujita and Florentin Smarandache. A reconsideration of advanced concepts in neutrosophic graphs: Smart, zero divisor, layered, weak, semi, and chemical graphs. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 308.
 - [53] Takaaki Fujita and Florentin Smarandache. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond (Second Volume)*. Biblio Publishing, 2024.
 - [54] Takaaki Fujita and Florentin Smarandache. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond (Third Volume)*. Biblio Publishing, 2024.
 - [55] Takaaki Fujita and Florentin Smarandache. Antipodal turiyam neutrosophic graphs. *Neutrosophic Optimization and Intelligent Systems*, 5:1–13, 2024.
 - [56] Takaaki Fujita and Florentin Smarandache. Introduction to upside-down logic: Its deep relation to neutrosophic logic and applications. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond (Third Volume)*, 2024.
 - [57] Takaaki Fujita and Florentin Smarandache. Mixed graph in fuzzy, neutrosophic, and plithogenic graphs. *Neutrosophic Sets and Systems*, 74:457–479, 2024.
 - [58] Takaaki Fujita and Florentin Smarandache. A review of the hierarchy of plithogenic, neutrosophic, and fuzzy graphs: Survey and applications. In *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond (Second Volume)*. Biblio Publishing, 2024.
 - [59] Takaaki Fujita and Florentin Smarandache. A short note for hypersoft rough graphs. *HyperSoft Set Methods in Engineering*, 3:1–25, 2024.
 - [60] Takaaki Fujita and Florentin Smarandache. Study for general plithogenic soft expert graphs. *Plithogenic Logic and Computation*, 2:107–121, 2024.
 - [61] Takaaki Fujita and Florentin Smarandache. Uncertain automata and uncertain graph grammar. *Neutrosophic Sets and Systems*, 74:128–191, 2024.
 - [62] Takaaki Fujita and Florentin Smarandache. Local-neutrosophic logic and local-neutrosophic sets: Incorporating locality with applications. *Advancing Uncertain Combinatorics through Graphization, Hyperization, and Uncertainization: Fuzzy, Neutrosophic, Soft, Rough, and Beyond*, page 51, 2025.
 - [63] Harish Garg, Muhammad Riaz, Muhammad Abdullah Khokhar, and Maryam Saba. Correlation measures for cubic m-polar fuzzy sets with applications. *Mathematical Problems in Engineering*, 2021.
 - [64] W-L Gau and Daniel J Buehrer. Vague sets. *IEEE transactions on systems, man, and cybernetics*, 23(2):610–614, 1993.
 - [65] Daniela Gifu. Soft sets extensions: Innovating healthcare claims analysis. *Applied Sciences*, 14(19):8799, 2024.
 - [66] Muhammad Gulistan, Naveed Yaqoob, Ahmed Elmoasry, and Jawdat Alebraheem. Complex bipolar fuzzy sets: An application in a transport’s company. *J. Intell. Fuzzy Syst.*, 40:3981–3997, 2021.
 - [67] Masooma Raza Hashmi, Muhammad Riaz, and Florentin Smarandache. m-polar neutrosophic topology with applications to multi-criteria decision-making in medical diagnosis and clustering analysis. *International Journal of Fuzzy Systems*, 22:273–292, 2020.
 - [68] Karel Hrbacek and Thomas Jech. Introduction to set theory, revised and expanded. 2017.
 - [69] Satham Hussain, Jahir Hussain, Isnaini Rosyida, and Said Broumi. Quadripartitioned neutrosophic soft graphs. In *Handbook of Research on Advances and Applications of Fuzzy Sets and Logic*, pages 771–795. IGI Global, 2022.
 - [70] Satham S Hussain, Muhammad Aslam, Hossein Rahmonlou, and N Durga. Applying interval quadripartitioned single-valued neutrosophic sets to graphs and climatic analysis. In *Data-Driven Modelling with Fuzzy Sets*, pages 100–143. CRC Press, 2025.
 - [71] Thomas Jech. *Set theory: The third millennium edition, revised and expanded*. Springer, 2003.
 - [72] Kazimierz Kuratowski. Introduction to set theory and topology. 1964.
 - [73] An Lu and Wilfred Ng. Vague sets or intuitionistic fuzzy sets for handling vague data: which one is better? In *International conference on conceptual modeling*, pages 401–416. Springer, 2005.
 - [74] Pradip Kumar Maji, Ranjit Biswas, and A Ranjan Roy. Soft set theory. *Computers & mathematics with applications*, 45(4-5):555–562, 2003.
 - [75] M. A. Malik. m-polar neutrosophic graphs. 2021.
 - [76] Rama Mallick and Surapati Pramanik. Pentapartitioned neutrosophic set and its properties. *Neutrosophic Sets and Systems*, 35:49, 2020.
 - [77] Mai Mohamed and Asmaa Elsayed. A novel multi-criteria decision making approach based on bipolar neutrosophic set for evaluating financial markets in egypt. *Multicriteria Algorithms with Applications*, 2024.
 - [78] Kartick Mohanta, Arindam Dey, Anita Pal, Hoang Viet Long, and Le Hoang Son. A study of m-polar neutrosophic graph with applications. *Journal of Intelligent & Fuzzy Systems*, 38(4):4809–4828, 2020.
 - [79] Dmitriy Molodtsov. Soft set theory-first results. *Computers & mathematics with applications*, 37(4-5):19–31, 1999.

-
- [80] T Mythili, V Jeyanthi, and KR Senthilkumar. Enhanced decision-making academic libraries with topsis methods using blockchain technology by heptapartitioned neutrosophic number. In *Leveraging Blockchain for Future-Ready Libraries*, pages 123–140. IGI Global Scientific Publishing, 2025.
 - [81] M Myvizhi, Ahmed M Ali, Ahmed Abdelhafeez, and Haitham Rizk Fadlallah. *MADM Strategy Application of Bipolar Single Valued Heptapartitioned Neutrosophic Set*. Infinite Study, 2023.
 - [82] M Myvizhi, Mohamed Elkholy, Ahmed Abdelhafeez, Hussam Elbehieri, et al. Enhanced madm strategy with heptapartitioned neutrosophic distance metrics. *Neutrosophic Sets and Systems*, 78:74–96, 2025.
 - [83] Khalid Naeem and Bijan Divvaz. Information measures for madm under m-polar neutrosophic environment. *Granular Computing*, 8(3):597–616, 2023.
 - [84] Junda Qiu, Linjia Jiang, Honghui Fan, Peng Li, and Congzhe You. Dynamic nonlinear simplified neutrosophic sets for multiple-attribute group decision making. *Heliyon*, 10(5), 2024.
 - [85] Radha R. and Stanis Arul Mary. Pentapartitioned neutrosophic pythagorean set. *International Research Journal on Advanced Science Hub*, 2021.
 - [86] R. Radha, A. Stanis Arul Mary, and Florentin Smarandache. Quadripartitioned neutrosophic pythagorean soft set. *International Journal of Neutrosophic Science*, 2021.
 - [87] Ramakrishnan Radha, A. Stanis, and Arul Mary. Pentapartitioned neutrosophic pythagorean soft set. 2021.
 - [88] M Ramya, Sandesh Murali, and R.Radha. Bipolar quadripartitioned neutrosophic soft set. 2022.
 - [89] M Murali Krishna Rao. Tripolar fuzzy interior ideals and tripolar fuzzy soft interior ideals over semigroups. *Annals of Fuzzy Mathematics and Informatics*, 20(3):243–256, 2020.
 - [90] M Murali Krishna Rao and B Venkateswarlu. Tripolar fuzzy interior ideals of a γ -semiring. *Asia Pacific Journal of Management*, 5(2):192–207, 2018.
 - [91] M Murali Krishna Rao, B Venkateswarlu, and Y Adi Narayana. Tripolar fuzzy soft ideals and tripolar fuzzy soft interior ideals over semiring. *Italian journal of pure and applied Mathematics*, (422019):731743, 2019.
 - [92] Muhammad Riaz, Muhammad Tahir Hamid, Deeba Afzal, Dragan Pamucar, and Yuming Chu. Multi-criteria decision making in robotic agri-farming with q-rung orthopair m-polar fuzzy sets. *PLoS ONE*, 16, 2021.
 - [93] Muhammad Riaz and Masooma Raza Hashmi. Soft rough pythagorean m-polar fuzzy sets and pythagorean m-polar fuzzy soft rough sets with application to decision-making. *Computational and Applied Mathematics*, 39, 2019.
 - [94] Judith Roitman. *Introduction to modern set theory*, volume 8. John Wiley & Sons, 1990.
 - [95] Muhammad Saeed, Muhammad Saqlain, and Asad Mehmood. *Application of similarity measure on m-polar interval-valued neutrosophic set in decision making in sports*, volume 38. Infinite Study, 2020.
 - [96] AA Salama, KF Alhasan, HA Elagamy, and Florentin Smarandache. Neutrosophic dynamic set. *Neutrosophic Knowledge*, 3, 2021.
 - [97] Sovan Samanta, Madhumangal Pal, Hossein Rashmanlou, and Rajab Ali Borzooei. Vague graphs and strengths. *Journal of Intelligent & Fuzzy Systems*, 30(6):3675–3680, 2016.
 - [98] Muhammad Saqlain and Xiao Long Xin. *Interval valued, m-polar and m-polar interval valued neutrosophic hypersoft sets*. Infinite Study, 2020.
 - [99] S Satham Hussain, Durga Nagarajan, Hossein Rashmanlou, and Farshid Mofidnakhai. Novel supply chain decision making model under m-polar quadripartitioned neutrosophic environment. *Journal of Applied Mathematics and Computing*, pages 1–26, 2024.
 - [100] Xiaolong Shi, Saeed Kosari, Hossein Rashmanlou, Said Broumi, and S Satham Hussain. Properties of interval-valued quadripartitioned neutrosophic graphs with real-life application. *Journal of Intelligent & Fuzzy Systems*, 44(5):7683–7697, 2023.
 - [101] Bimal Shil, Rakhal Das, Carlos Granados, Suman Das, and Binod Chandra Tripathy. Single-valued quadripartitioned neutrosophic d-ideal of d-algebra. *Neutrosophic Sets and Systems*, 67:105–114, 2024.
 - [102] M Sivakumar, Abdul Rajak Rabiyyathul Basariya, M Senthil, G Raja, R Rajavarman, et al. Transforming arabic text analysis: Integrating applied linguistics with m-polar neutrosophic set mood change and depression on social media. *International Journal of Neutrosophic Science*, (2):313–13, 2025.
 - [103] F. Smarandache. Introduction to superhyperalgebra and neutrosophic superhyperalgebra. *Journal of Algebraic Hyperstructures and Logical Algebras*, 2022.
 - [104] Florentin Smarandache. Ambiguous set (as) is a particular case of the quadripartitioned neutrosophic set (qns). *nidus idearum*, page 16.
 - [105] Florentin Smarandache. Neutrosophic overset, neutrosophic underset, and neutrosophic offset. similarly for neutrosophic over-/under-/offlogic, probability, and statistics neutrosophic, pons editions brussels, 170 pages book, 2016.
 - [106] Florentin Smarandache. Neutrosophy: neutrosophic probability, set, and logic: analytic synthesis & synthetic analysis. 1998.
 - [107] Florentin Smarandache. A unifying field in logics: Neutrosophic logic. In *Philosophy*, pages 1–141. American Research Press, 1999.
 - [108] Florentin Smarandache. Hyperuncertain, superuncertain, and superhyperuncertain sets/logics/probabilities/statistics. *Critical Review*, XIV, 2017.
 - [109] Florentin Smarandache. *Plithogeny, plithogenic set, logic, probability, and statistics*. Infinite Study, 2017.
 - [110] Florentin Smarandache. Extension of soft set to hypersoft set, and then to plithogenic hypersoft set. *Neutrosophic sets and systems*, 22(1):168–170, 2018.
 - [111] Florentin Smarandache. *Plithogenic set, an extension of crisp, fuzzy, intuitionistic fuzzy, and neutrosophic sets-revisited*. Infinite study, 2018.
 - [112] Florentin Smarandache. Plithogeny, plithogenic set, logic, probability, and statistics. *arXiv preprint arXiv:1808.03948*, 2018.

-
- [113] Florentin Smarandache. Foundation of superhyperstructure & neutrosophic superhyperstructure. *Neutrosophic Sets and Systems*, 63(1):21, 2024.
- [114] Florentin Smarandache and NM Gallup. Generalization of the intuitionistic fuzzy set to the neutrosophic set. In *International Conference on Granular Computing*, pages 8–42. Citeseer, 2006.
- [115] Nguyen Tho Thong, Luong Thi Hong Lan, Shuo-Yan Chou, Le Hoang Son, Do Duc Dong, and Tran Thi Ngan. An extended topsis method with unknown weight information in dynamic neutrosophic environment. *Mathematics*, 8(3):401, 2020.
- [116] Nguyen Tho Thong, Luu Quoc Dat, Nguyen Dinh Hoa, Mumtaz Ali, Florentin Smarandache, et al. Dynamic interval valued neutrosophic set: Modeling decision making in dynamic environments. *Computers in Industry*, 108:45–52, 2019.
- [117] Nguyen Tho Thong, Florentin Smarandache, Nguyen Dinh Hoa, Le Hoang Son, Luong Thi Hong Lan, Cu Nguyen Giap, Dao The Son, and Hoang Viet Long. A novel dynamic multi-criteria decision making method based on generalized dynamic interval-valued neutrosophic set. *Symmetry*, 12(4):618, 2020.
- [118] Vakkas Ulucay, Irfan Deli, and Mehmet Sahin. Similarity measures of bipolar neutrosophic sets and their application to multiple criteria decision making. *Neural Computing and Applications*, 29:739–748, 2018.
- [119] Jun Ye. Correlation coefficient between dynamic single valued neutrosophic multisets and its multiple attribute decision-making method. *Information*, 8(2):41, 2017.
- [120] Lotfi A Zadeh. Fuzzy sets. *Information and control*, 8(3):338–353, 1965.
- [121] Lotfi A Zadeh. A fuzzy-set-theoretic interpretation of linguistic hedges. 1972.
- [122] Lotfi A Zadeh. Fuzzy sets and their application to pattern classification and clustering analysis. In *Classification and clustering*, pages 251–299. Elsevier, 1977.
- [123] Lotfi A Zadeh. Fuzzy sets versus probability. *Proceedings of the IEEE*, 68(3):421–421, 1980.
- [124] Lotfi A Zadeh. Fuzzy logic, neural networks, and soft computing. In *Fuzzy sets, fuzzy logic, and fuzzy systems: selected papers by Lotfi A Zadeh*, pages 775–782. World Scientific, 1996.